

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.



wherein:

A is a member selected from the group consisting of: R^2 , $-NR^3R^4$, $-C(=O)NR^3R^4$,



where R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system having 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the group consisting of N, O and S;

m is an integer from 0-3;

Z is a member selected from the group consisting of a direct link, C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, or a five to ten membered

19 heterocyclic ring system having 1-4 heteroatoms selected from the group consisting of N,
20 O and S;

21 n is an integer from 0-3;

22 D is a member selected from the group consisting of a direct link, -CH₂-, -O-,
23 -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-,
24 -C(=O)-N(R²)- and -N(R²)-C(=O)-;

25 R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋
26 ₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH,
27 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³,
28 -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino
29 group, wherein the substituted amino groups are independently substituted by at least one
30 member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl,
31 C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and
32 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, where R² and R³ is as described above;

33 q is an integer from 0-3;

34 R¹¹ and R¹² are independently a member selected from the group consisting of H,
35 C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl,
36 C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰,
37 -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰,
38 -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and
39 R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋
40 ₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
41 form a saturated or unsaturated ring with the atom to which they are both attached;

42 p is an integer from 0-3;


43 E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)-,
44 where R¹¹ is as set forth above, phenylene, a bivalent 5 to 12 member heteroaryl group
45 having 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to
46 ten membered non-aromatic bivalent heterocyclic ring system having 1-4 heteroatoms

a'
47 selected from the group consisting of N, O and S, wherein said heteroaryl and said non-
48 aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴
49 groups;

50 J is a member selected from the group consisting of a direct link, a bivalent
51 C₃₋₈cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group having 1 to
52 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten
53 membered non-aromatic bivalent heterocyclic ring system having 1-4 heteroatoms
54 selected from the group consisting of N, O and S wherein said heteroaryl and said non-
55 aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴
56 groups;

57 each R¹⁴ group is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈-
58 alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH,
59 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkyl-OH, C₀₋₈alkyl-SH, -O-R² and
60 -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group,
61 wherein the substituted amino groups are independently substituted by at least one
62 member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl,
63 C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

64 G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



65 wherein


66 t is an integer from 0 to 6,

67 u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are
68 independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring
69 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
70 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
71 such atoms being selected from the group consisting of N, O and S; where R^{18} taken with
72 R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently
73 form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the
74 group consisting of N, O and S;

75
76 with the proviso that when G is H, -CN, -OR¹⁷, either E or J must contain at least
77 one N atom;

78 or a pharmaceutically acceptable diastereomer, salt, hydrate, and solvate thereof.


1 2. (Amended) A compound of formula II:



2

3

A is a member selected from the group consisting of:



4
 5 where R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H,
 6 -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to
 7 ten membered heterocyclic ring system having 1-4 heteroatoms selected from the group
 8 consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system
 9 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O
 10 and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a
 11 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the group
 12 consisting of N, O and S;

13 Z is a member selected from the group consisting of C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{2-8}
 14 alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, and a five to ten membered heterocyclic ring
 15 system having 1-4 heteroatoms selected from the group consisting of N, O and S;

16 D is a member selected from the group consisting of a direct link, -CH₂-, -O-,
 17 -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-,
 18 -C(=O)-N(R²)- and -N(R²)-C(=O)-, provided that when Z is C_{1-8} alkyl, C_{2-8} alkenyl,
 19 C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, then D is -O-, or -N(R²)-;

20 R¹ is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8}
 21 alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH,
 22 C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{1-8} alkyl-OH, C_{0-8} alkyl-SH, -C(=O)NR²R³,
 23 -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino

a!
24 group, wherein the substituted amino groups are independently substituted by at least one
25 member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl,
26 C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and
27 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;


28 R² and R³ are independently selected from the group consisting of H, -OH,
29 C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten
30 membered heterocyclic ring system having 1-4 heteroatoms selected from the group
31 consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system
32 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O
33 and S;

34 q is an integer from 0-3;

35 R¹¹ is independently a member selected from the group consisting of H, C₁₋₈alkyl,
36 C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl,
37 C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰,
38 -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰,
39 -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and
40 R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈
41 alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
42 form a saturated or unsaturated ring with the atom to which they are both attached;
43 p is an integer from 0-2;

44 E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)- ,
45 where R¹¹ is as set forth above, phenylene, a bivalent 5 to 12 member heteroaryl group
46 having 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to
47 ten membered non-aromatic bivalent heterocyclic ring system having 1-4 heteroatoms
48 selected from the group consisting of N, O and S, wherein said heteroaryl and said non-
49 aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴
50 groups;

65 G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



66 wherein

67 t is an integer from 0 to 6,

68 u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are
69 independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring
70 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
71 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
72 such atoms being selected from the group consisting of N, O and S; where R^{18} taken with
73 R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently
74 form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the
75 group consisting of N, O and S;


76 with the proviso that when G is H, -CN, -OR¹⁷, either E or J must contain at least
77 one N atom;

78 or a pharmaceutically acceptable diastereomer, salt, hydrate, and solvate thereof.

80

1 5. (Amended) A compound of formula III:

2



3

4 wherein:

5 R^8 is selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring
6 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
7 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
8 such atoms being selected from the group consisting of N, O and S;

9

Q²
C¹
cont

10 R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋
11 8alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH,
12 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³,
13 -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino
14 group, wherein the substituted amino groups are independently substituted by at least one
15 member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl,
16 C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and
17 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, where R² and R³ is as described above;

18 R² is selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋
19 8alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring
20 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
21 C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
22 such atoms being selected from the group consisting of N, O and S;
23 q is 0-3;

24 R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl,
25 C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl,
26 -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰,
27 -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰,
28 -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member
29 selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein
30 when two R¹⁰ groups are present they may be taken together to form a saturated or
31 unsaturated ring with the atom to which they are both attached;
32 p is an integer from 0-2;


33 E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)- ,
34 where R¹¹ is as set forth above, phenylene, a bivalent 5 to 12 member heteroaryl group
35 having 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to
36 ten membered non-aromatic bivalent heterocyclic ring system having 1-4 heteroatoms

37 selected from the group consisting of N, O and S, wherein said heteroaryl and said non-
38 aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴
39 groups;

40 J is a member selected from the group consisting of a direct link, a bivalent
41 C₃₋₈cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group having 1 to
42 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten
43 membered non-aromatic bivalent heterocyclic ring system having 1-4 heteroatoms
44 selected from the group consisting of N, O and S wherein said heteroaryl and said non-
45 aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴
46 groups;

47 each R¹⁴ group is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈-
48 alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH,
49 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₁₋₈alkyl-OH, C₀₋₈alkyl-SH, -O-R² and
50 -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group,
51 wherein the substituted amino groups are independently substituted by at least one
52 member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl,
53 C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

54 G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



55 wherein


56 t is an integer from 0 to 6,

57 u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are
58 independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring
59 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
60 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
61 such atoms being selected from the group consisting of N, O and S; where R^{18} taken with
62 R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently
63 form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the
64 group consisting of N, O and S;

65 with the proviso that when G is H, -CN, -OR¹⁷, either E or J must contain at least
66 one N atom;

67 or a pharmaceutically acceptable diastereomer, salt, hydrate, and solvate thereof.
68



1 9. A compound of formula IV:



2

3 wherein:

4 A is a member selected from the group consisting of: R^2 , $-NR^3R^4$, $-C(=O)NR^3R^4$,



5
6 where $\text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8$, and R^9 are independently selected from the group
7 consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic
8 aryl, a five to ten membered heterocyclic ring system having 1-4 heteroatoms selected
9 from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in
10 the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group
11 consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with
12 R^8 , can each form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected
13 from the group consisting of N, O and S;

14 Z is a member selected from the group consisting of a direct link, C_{1-8} alkyl,
15 C_{3-8} cycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, or a five to ten membered
16 heterocyclic ring system having 1-4 heteroatoms selected from the group consisting of N,
17 O and S;

18 n is 0-3;

19 D is a member selected from the group consisting of $-\text{CH}_2-$, $-\text{O}-$, $-\text{NR}^2$, $-\text{C}(=\text{O})-$,
20 $-\text{S}-$, $-\text{SO}_2-$, $-\text{SO}_2-\text{NR}^2$, $-\text{NR}^2-\text{SO}_2$, $-\text{OC}(=\text{O})-$, $-\text{C}(=\text{O})\text{NR}^2$, and $-\text{NR}^2-\text{C}(=\text{O})-$;


21 R^1 and R^{14} are independently a member selected from the group consisting of H,
22 C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl,
23 C_{0-8} alkyl- $\text{C}(=\text{O})\text{OH}$, C_{0-8} alkyl- $\text{C}(=\text{O})\text{O}-\text{C}_{1-8}$ alkyl, $-\text{CN}$, $-\text{NO}_2$, C_{1-8} alkyl-OH,
24 C_{0-8} alkyl-SH, $-\text{O}-\text{R}^2$ and $-\text{O}-\text{C}(=\text{O})\text{R}^2$, an unsubstituted amino group, a mono- or
25 di-substituted amino group, wherein the substituted amino groups are independently

26 substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋
 27 ₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and
 28 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

29 q is 0-3;

30 R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl,
 31 C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl,
 32 -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰,
 33 -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰,
 34 -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member
 35 selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein
 36 when two R¹⁰ groups are present they may be taken together to form a saturated or
 37 unsaturated ring with the atom to which they are both attached;

38 G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



39 wherein

40 t is an integer from 0 to 6,


41 u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are

a³
c¹
cont

42 independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋
43 galkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring
44 system having 1-4 heteroatoms selected from the group consisting of N, O and S; and
45 C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of
46 such atoms being selected from the group consisting of N, O and S; where R¹⁸ taken with
47 R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently
48 form a 5 to 6 membered heterocyclic ring having from 1 to 4 atoms selected from the
49 group consisting of N, O and S;
50 with the proviso that when G is H, -CN, -OR¹⁷, either E or J must contain at least
51 one N atom;
52 or a pharmaceutically acceptable diastereomer, salt, hydrate, and solvate thereof.

a⁴

1 11. A compound of formula V:



2
3 wherein:

4 R², R⁶, and R⁹ are independently selected from the group consisting of H, -OH,
5 C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten
6 membered heterocyclic ring system having 1-4 heteroatoms selected from the group
7 consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system
8 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O
9 and S;

10 R¹¹ is independently a member selected from the group consisting of H,
11 C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl,